

Potential CO₂ Geological Storage Sites for Carbon Capture and Storage (CCS) in Québec St. Lawrence Lowlands - A Preliminary Analysis

K. Bédard*, M. Malo

Institut National de la recherche scientifique, Québec, QC, Canada
karine.bedard@ete.inrs.ca

Y. Duchaine

Consultant, Trois-Rivières, QC, Canada

E. Konstantinovskaya, B. Giroux

Institut National de la recherche scientifique, Québec, QC, Canada

Summary

The St. Lawrence Lowlands platform and frontal thrust slices of the Appalachian belt are prospective for CO₂ storage in the province of Québec. The thick (1500 to 3000 meters) sedimentary succession hosts two types of reservoirs: deep saline aquifers in the platform and gas reservoirs in the fold-thrust belt. The Potsdam, Beekmantown and Trenton groups are identified as the potential reservoir formations in the St. Lawrence Lowlands sedimentary succession. A preliminary analysis of selected storage sites is used to evaluate theoretical and effective capacities for CO₂ storage at the basin and reservoir scales.

Introduction

Interest for CCS is rapidly growing as it represents a short-to-medium-term solution for reducing anthropogenic CO₂ emissions in the atmosphere (IEA, 2008; IPCC, 2005). The technology for storing CO₂ in oil and gas reservoirs and in deep saline aquifers is now ready to be applied in the context of CCS (Bachu, 2008). Research for potential geological storage sites is thus going on worldwide where there are large sedimentary basins known as hosting geological reservoirs.

The objectives of the research project are to determine the potential capacity of CO₂ geological storage in Québec in order to evaluate the feasibility of CCS in the province. The St. Lawrence Lowlands is the current prospect area for CO₂ storage in the south part of Québec province where most of the large CO₂ emitters are located. The most advanced methodologies are used to identify the prospective CO₂ geological storage sites in Québec.

Geological setting

The Cambro-Ordovician platform sedimentary succession of the St. Lawrence Lowlands has a thickness of 1500 to 3000 meters and was unconformably deposited on the Precambrian bedrock (Globensky, 1987). The sedimentary succession is composed of sandstones of the Potsdam Group, dolomites of the Beekmantown Group and limestones of the Trenton, Black River and Chazy groups. They are overlain by the Utica Shale and the siltstones and shales of the Lorraine Group. SW-NE normal faults affect the succession which deepens toward the south-east (Castonguay *et al.*, 2006). The Appalachian fold-thrust belt is limited by the Aston Fault and the Logan's Line, in between which we find parautochthonous parts of the platform sedimentary succession caught up in fault slices and displaced above rocks of the platform (figures 1 and 2).




Figure 1: Southern Québec structural domains and location of the M-2001 seismic line (figure 2) (modified from Bertrand *et al.*, 2003).

Deep saline aquifers are found in the sedimentary succession of the platform, particularly in the Potsdam, Beekmantown and Trenton groups. The gas reservoirs, on the other hand, are known to be located in imbricated frontal thrust slices of the Appalachian fold-thrust belt (e.g. the Saint-Flavien reservoir, figure 2).




Figure 2: Architecture of the St. Lawrence Lowlands platform and the frontal part of the Appalachian fold-thrust belt, the two domains hosting potential CO₂ storage sites, on seismic line M-2001 (modified from Castonguay *et al.*, 2006).

Criteria for assessing CO₂ storage sites

As not all the sedimentary basins are appropriate for CO₂ sequestration, a preliminary analysis must be conducted to evaluate their potential. Many geological and practical criteria must be taken into account in this assessment. Some of these criteria include depth, size, tectonic setting of the basin, fault occurrences, presence of an extensive vertical and horizontal seal (caprock), accessibility, infrastructures and proximity of large CO₂ sources (Bachu, 2003).

At the reservoir scale, the evaluation is more specific. The positive indicators include, among others, a depth of 1000 m to 2500 m, a reservoir thickness of at least 50 m, at least 20% of porosity and 300 mD of permeability, an unfaulted continuous caprock with a thickness of at least 100 m (Chadwick *et al.*, 2008).

In the case of the St. Lawrence Lowlands, a preliminary evaluation conducted by the chair on geological CO₂ sequestration at INRS is used to determine the potential of the basin and the reservoirs of the St. Lawrence Lowlands.

Preliminary analysis of the St. Lawrence Lowlands

The analysis of the criteria for assessing the potential of the St. Lawrence Lowlands basin shows many positive indicators for CO₂ geological sequestration. First of all, the basin extends over more than 10 000 km² which is a fairly good size and, most importantly, it is covered by a extensive caprock that can exceed 1000 m of thickness (the Utica Shale and the Lorraine Group). The St. Lawrence Lowlands sedimentary units that may represent potential reservoir capacities for the CO₂ storage are located in sandstones of the Potsdam Group, dolomites of the Beekmantown Group, and more or less argillaceous limestones of the Trenton Group (including the thin Black River and Chazy groups). These sedimentary successions are found at depths of 1000 m to 2500 m within the platform and the imbricated thrust slices of the fold-thrust belt. The Potsdam Group is up to 750 m thick, its porosity is usually 6-12% and its permeability can be more than 10 mD and up to 200 mD in some thin layers. The Beekmantown dolomites are up to 450 m thick, they show porosities of 3-12%, exceptionally up to 18%, and good permeabilities varying from 45 mD to more than 250 mD. Finally, the Chazy-Black River-Trenton unit has a thickness of up to 375 m, lower porosities of 3-10% and permeability generally of the order of 1 mD but which can be more than 300 mD where saline aquifers are present.

Many practical criteria are positively evaluated such as: ease of access of the region, well developed infrastructures and several large CO₂ emitters located directly in the St. Lawrence Lowlands. Even if the porosities and permeabilities are lower than the positive indicators of Chadwick *et al.* (2008), the other good parameters still allow considering the St. Lawrence Lowlands units as potential CO₂ sequestration reservoirs.

Conclusions

The potential for CO₂ sequestration is evaluated in deep saline aquifers of the St. Lawrence Lowlands platform as well as in the gas reservoirs of the frontal thrust slices of the Appalachian fold-thrust belt. The effective capacity of the St. Lawrence Lowlands is assessed at the basin-scale (Bachu *et al.*, 2007) with the ultimate objective to identify specific sequestration sites and evaluate their practical and matched capacities for CO₂ sequestration (local-scale and site-scale assessment; Bachu *et al.*, 2007).

Acknowledgements

We would like to thank the “Ministère du Développement durable, de l’Environnement et des Parc du Québec” for the financing of the research chair of geological CO₂ sequestration. We also thank JUNEX for sharing data and knowledge with us as well as Mathieu Duchesne from the Geological Survey of Canada for reviewing initial versions of this contribution.

References

- Bachu, S., 2003. Screening and ranking of sedimentary basins for sequestration of CO₂ in geological media in response to climate change. *Environmental Geology*, 44, 277-289.
- Bachu, S., 2008. Legal and regulatory challenges in the implementation of CO₂ geological storage: An Alberta and Canadian perspective. *International Journal of Greenhouse Gas Control*, 2, 259-273.
- Bachu, S., Bonijoly, D., Bradshaw, J., Burruss, R., Holloway, S., Christensen, N. P., Maathiasen, O, M., 2007. CO₂ storage capacity estimation: Methodology and gaps. *International Journal of Greenhouse Gas Control*, 1, 430-443.
- Bertrand, R., Chagnon, A., Duchaine, Y., Lavoie, D., Malo, M., Savard, M. M., 2003. Sedimentologic, diagenetic and tectonic evolution of the Saint-Flavien gas reservoir at the structural front of the Québec Appalachians. *Bulletin of Canadian Petroleum Geology*, 51, 126–154.
- Castonguay, S., Dietrich, J., Shinduke, R., Laliberté, J-Y., 2006. Nouveau regard sur l'architecture de la Plate-forme du Saint-Laurent et des Appalaches du sud du Québec par le retraitement des profils de sismique réflexion M-2001, M-2002 et M-2003. Geological Survey of Canada, Open File 5328, 19 pages (2 sheets).
- Chadwick, A., Arts, R., Bernstone, C., May, F., Thibeau, S., Zweigel, P., 2008. Best Practice for the CO₂ Storage in Saline Aquifers – Observations and Guidelines from the SACS and CO2STORE Projects. British Geological Survey, Nottingham, UK, 273 pages.
- Globensky, Y., 1987. Géologie des Basses-Terres du Saint-Laurent. Ministère de l'Énergie et des Ressources du Québec, MM 85-02, 63 pages.
- Intergovernmental Panel on Climate Change (IPCC), 2005. IPCC Special Report on Carbon Dioxide Capture and Storage. IPCC, Cambridge University Press, 442 pages.
- International Energy Agency (IEA), 2008. CO₂ capture and storage - A key carbon abatement option. IEA/OECD, Paris, France, 261 pages.