

# The Saglek Basin in the Labrador Sea; past exploration history, current estimates and future opportunities

Christopher D. Jauer, Geological Survey of Canada  
cjauer@nrcan.gc.ca

and

Gordon N. Oakey<sup>1</sup>, Mike Avery<sup>1</sup>, Graham Williams<sup>1</sup> and Hans Wielens<sup>\*1</sup>

<sup>1</sup> Geological Survey of Canada ( Atlantic ), 1 Challenger Drive, Dartmouth, Nova Scotia, Canada, B2Y 4A2

\* deceased

## GeoConvention 2012: Vision

New interpretations of industry seismic data in northeast Labrador Sea have been produced to develop a 3-D Petroleum System Model for the Saglek Basin. Improved palynological analysis has provided better age constraints on the timing and geometry of basin in-filling and subsidence. Exploration well cutting samples have been re-analyzed to improve the details of thermal maturity and kerogen source type. A 2-component heat-flow model has been adopted based on the tectonic history of the Labrador Sea and the timing of significant volcanic extrusions each producing significant heat pulses. The new petroleum model explains the origins of the discovery at Hekja 0-71 and provides an explanation for the dry Raleigh-N18 well only 25 km away. The revised seismic mapping used for the model building shows the presence of very large structural closures that could represent significant future exploration targets; one very large structure mapped is over 10 times the size of the Hekja structure.

The Saglek Basin is a very large sedimentary basin (over 100,000 km<sup>2</sup>) on the northern end of the Labrador Margin. The 2 Tcf Hekja 0-71 discovery in 1979 lies at the northern end of the Saglek Basin near the mouth of Frobisher Bay offshore Nunavut. The southern half of the basin is within the offshore jurisdiction of Newfoundland & Labrador. The basin contains over 9 km of strata mapped by regional 2-D seismic data.

A basin-wide seismic reflector that is interpreted as basalts in the north leads into a regional unconformity in the south. These basalts have been sampled from the exploration wells, and are believed to be part of a major subareal Paleocene volcanic event corresponding with the initiation of sea-floor spreading (Chron 27N).

A second volcanic episode corresponded with a major change in direction of motion of the Greenland Plate (Chron 24R). Satellite-derived gravity data (Sandwell and Smith, 2009) have been used to produce a marine Bouguer gravity anomaly map, which shows that the now-extinct (Chron 13N) Labrador Sea spreading axis extends into the Saglek Basin and cannot be seismically differentiated from the mapped "basalt" seismic horizon. Based on these tectonically controlled volcanic events, two heat pulses at 65Ma and 56Ma were used in the basin thermal history used in the petroleum system model.

The geochemical analysis of the well cuttings from the Hekja 0-71 well indicates that they were thermally too immature to produce the gas and condensate found at the depth of discovery, ~3200 m (Fowler et al., 2005). Terrestrial-based source rocks (deltaic siliciclastics with coals) containing resinates are capable of generating hydrocarbons at low maturity levels (Snowden, 1991); however the model suggests that these hydrocarbons were not locally "self-sourced" but filled the structural trap by

fluid migration from a deeper more thermally mature deposit of similar source rocks. We believe that Upper Cretaceous strata beneath the Paleocene basalts were also a significant contributor to the petroleum generation in the basin and the model results show that original in-place gas volumes of 170 to 260 Tcf are possible.

A new synthesis of the exploration well stratigraphic correlations was made possible by new palynological analysis that greatly improved age dating and depositional environments. Basement blocks responded to changes in the regional tectonic episode, forming numerous structural traps of which only a few were drilled. The basement controlled faults created sediment bypass fairways in some areas resulting in a complex distribution of coastal siliciclastic facies further seaward than expected. Three prospective formations capable of generating petroleum have been identified, and their variable depth-of-burial and proximity to the siliciclastic reservoir rocks are critical factors in the modelling of petroleum generation and trapping.

Although all major tectonic movements ceased in the late Eocene (Chron 13N, ~35 Ma) the area is still seismically active. The "Labrador Sea Seismic Zone" is a linear belt of medium-amplitude earthquakes coincident with the extinct spreading axis. Seismic events from offshore California are known to trigger venting of petroleum (Field and Jennings, 1987). In the Saglek Basin, there is an unaccounted lack of observed reflection seismic indicators of gas charged sediments or gas hydrates. The presence of seafloor features that appear to be caused by petroleum venting and oil slicks on the sea surface imaged by satellite radar (Jauer and Budkewitsch, 2009) suggests that there has been ongoing periodic seepage of petroleum from the basin and that this process may be augmented by persistent local earthquake triggers.

The existing understanding of petroleum potential for the Saglek Basin is based on limited seismic mapping, eight drilled exploration wells and one significant discovery. The petroleum system, previously believed to be locally derived and "self-sourcing", is now shown to be generated at greater depths and has migrated into the reservoirs over wide distances. The many untested structures and the hydrocarbon volumes generated by the model suggest that there is significant petroleum potential for the region


## References

Field, Micheal and Jennings, M.A., 1987, Seafloor gas seeps triggered by a northern California earthquake, *Marine Geology*, vol 77, issue 1-2, pp 39-51


Fowler, M., Stasiuk, L. and Avery, M., 2005 Potential petroleum systems in the Labrador and Baffin Shelf areas, offshore north eastern Canada. In *Organic Geochemistry: Challenges for the 21<sup>st</sup> Century Vol. 1. Abstracts for 22<sup>nd</sup> International Meeting on Organic Geochemistry*, Seville, Spain, September 2005 pg. 463-463

Sandwell, D. T., and W. H. F. Smith, 2009, Global marine gravity from retracked Geosat and ERS-1 altimetry: Ridge segmentation versus spreading rate, *J. Geophys. Res.*, 114, B01411, doi:10.1029/2008JB006008


Snowdon, Lloyd R., 1991. Oil from Type III organic matter: resinite revisited. *Organic Geochemistry*, Vol. 17, No. 6 pp. 743-747



## Seismic depth mapping of Gudrid Member at Hekja O-71 and Raleigh N-18 locations



from Jauer, 2009



## Acknowledgements

We wish to extend our thanks to support from Husky Oil and Suncor Energy for data access.